

CAUTION

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



ACTIVE CONSTITUENT: 750 g/kg MANCOZEB

GROUP M3 FUNGICIDE

**For the control of certain fungus diseases of
fruit, field crops, tobacco, turf, vegetables and ornamentals
as per Directions for Use table.**

GENERAL INSTRUCTIONS

- Fortuna Globe 750 WG is a protectant fungicide for the control of certain fungus diseases in many crops.
- For best results, apply as a regular spray program.
- Thorough coverage of the treated crop is essential.

FUNGICIDE RESISTANCE WARNING:



For fungicide resistance management Fortuna Globe 750 WG is a Group M3 fungicide.

Some naturally occurring individual fungi resistant to Fortuna Globe 750 WG and other Group M3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. The resistant fungi will not be controlled by Fortuna Globe 750 WG and other Group M3 fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, Grochem Australia Pty Ltd accepts no liability for any losses that may result from the failure of Fortuna Globe 750 WG to control resistant fungi.

MIXING

- Slowly place into tank as it is being filled, or thoroughly premix in a nurse tank for concentrate or aircraft sprayers. Add other co-applied fungicides, insecticides, surfactants etc after Fortuna Globe 750 WG has been placed into suspension.
- Maintain sufficient agitation during mixing and spraying to prevent the product settling out and to ensure a uniform spray mixture. Rinse out spray tank, pumps and nozzles at the end of the day.
- When preparing spray solutions for use in hand sprayer, premix as a slurry in a small container, and then add to sprayer containing one-third to half desired final water volume.

SURFACTANTS

- The addition of agricultural surfactants to Fortuna Globe 750 WG will improve initial spray deposits, fungicide redistribution and weatherability.
- Use a Spreader-Sticker per label directions if needed.

APPLICATION

Dilute Spraying – TREE AND VINE CROPS ONLY

- Use a sprayer designed to apply high volumes of water up to the point of run-off and match to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of first run-off. Avoid excessive run-off.
- The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from Industry guidelines or expert advice.
- Add the amount of product specified in the Directions for Use table for each 100 L of water. Spray to the point of run-off.
- The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

Concentrate Spraying – TREE AND VINE CROPS ONLY

- Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.
- Determine an appropriate dilute spray volume (see Dilute Spraying above) for the crop canopy. Consult your local advisor, agronomist or Department of Agriculture to determine this volume. This is needed to calculate the concentrate-mixing rate.
- The mixing rate for concentrate spraying can then be calculated in the following way.

Concentrate Spraying Example

1. Dilute spray volume as determined above: e.g. 1,000 L/ha;
 2. Your chosen concentrate spray volume: e.g. 500 L/ha;
 3. The concentration factor is $2 \times (1,000 \div 500)$, and;
 4. If the dilute label rate is 150 g/100 L, then the concentrate rate becomes 2×150 , i.e. 300 g/100 L of concentrate spray.
- The chosen spray volume, amount of product per 100 L of water and the sprayer set up and operation may need to be changed as the crop grows.
 - For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow Industry Best Practices.
 - For concentrate application, use a spray volume of 20 L or more per hectare.
 - For dilute application, apply to run-off. See Dilute Spraying above.

AERIAL APPLICATION

- May be applied by agricultural aircraft.
- Use at least 30 to 50 litres of spray mixture per hectare. Consult the Directions for Use for the correct rate to be applied in specific crop situations.

SPRAY TIMING

- Treatments should begin prior to disease infection and continue until the threat of disease has passed.
- Repeated applications are necessary to protect new plant growth.
- Reduce the spray interval when weather conditions favour disease development.

COMPATIBILITY

- Fortuna Globe 750 WG may be harmful to *Typhlodromus pyri*, and its use in orchards where integrated control is practiced should be avoided.
- This product is compatible with most commonly used agricultural fungicides and insecticides, however before preparing tank mixes, the user is advised to check compatibility with the other manufacturer's recommendations.

PROTECTION ON WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT:

- This product is toxic to fish
- Drift and run-off from treated areas may be hazardous to aquatic organisms in neighbouring areas.
- DO NOT contaminate ponds, waterways or drains with the chemical or used containers.
- DO NOT re-use container.

PROTECTION OF LIVESTOCK

- DO NOT graze treated turf or feed turf clippings from any treated area to poultry or livestock.

STORAGE AND DISPOSAL

- Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Keep away from fire and sparks.
- DO NOT allow to become wet or overheated in storage: decomposition, impaired activity or fire may result.
- DO NOT store near food, feedstuffs, fertilizers or seed. Users are reminded this product should be used before the stated expiry date.
- Shake empty bag into spray tank. Do not dispose of undiluted chemicals on site.
- Puncture or shred and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury empty packaging in a local authority landfill. If no landfill is available, bury the containers below 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations.
- Do not burn empty bags or product.

SAFETY DIRECTIONS

- May irritate the eyes, nose, throat and skin.
- Avoid contact with eyes and skin.
- DO NOT inhale dust.
- Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre, Phone Australia 131126. Avoid giving alcohol.

SAFETY DATA SHEET

If additional hazard information is required refer to the Safety Data Sheet.

For a copy phone 1800 777 068 or visit our website www.grochem.com

LIMIT OF LIABILITY

1. Grochem Australia accepts responsibility for the consistent quality of the product.
2. Grochem Australia accepts no responsibility whatsoever for any damage, injury or loss following purchase and use of this product.
3. The extent of liability of Grochem Australia is limited to the replacement of goods or a refund on the price paid. This being conditional upon a claim being made in writing and within 30 days of delivery/receipt of product.
4. This product must also be used in strict accordance with the directions as detailed on this label. The buyer accepts and uses this material with an understanding of the above conditions.

DIRECTIONS FOR USE

NON-TREE & VINE CROPS, FIELD CROP

CROP	DISEASES CONTROLLED	RATE/100 L	RATE/ha	WHP HARVEST (H) GRAZING (G)	CRITICAL COMMENTS
Cotton – Pima var. only	Alternaria leaf spot		2.5 kg	14 days (H)	Fortuna Globe 750 WG is a protective fungicide. To maximize control ensure thorough coverage of all plant surfaces. Begin applications as soon as disease symptoms appear. Repeat before each infection period (e.g. dew or rain) but no later than 7 to 10 days after the last application. DO NOT apply more than 4 sprays per season.
Fennel (Tas. only)	Leaf blight (<i>Cercosporidium</i>)	150-200g	2.2 kg		Apply as a ground spray if disease symptoms are present in November, allowing for one repeat treatment 10-14 days later. Aerial application may be required in prolonged wet periods.
Peanuts	Cercospora leaf spot, Rust		1.7 to 2.2 kg	14 days (H) 14 days (G)	Apply at 10 to 14 day intervals commencing when disease symptoms first appear. If rust is anticipated, apply at 7 to 10 day intervals. Use higher rate and shorter spray interval during weather conditions favourable to disease development.
Sunflowers For parent & hybrid seed production	Rust		2.2 kg	DO NOT GRAZE	Apply when disease symptom first appear and repeat at 7 to 10 day intervals while warm rainy weather conditions prevail. It is crucial to ensure application provides excellent coverage.
Poppies	Downy mildew		2.5 kg	7 weeks (H)	Apply as protective spray when weather conditions favour disease development. Repeat spray on 14-day intervals if conditions persist.
Tobacco	Blue mould Brown spot Frog eye leaf spot	200g	2.2 kg	Not required	Field treatments: Apply at 7 day intervals commencing 10 days after transplanting and ceasing 7 to 10 days before the first harvest. Increase the spray volume accordingly as plant size increases. Seed bed treatments: Apply when seedlings are 1.5 cm across and then repeat twice weekly. Wet the seedlings evenly without run-off.
Soybeans	Rust		2.2 kg	7 days (H) 14 days (G)	Apply when disease symptoms first appear and repeat at 7 to 10 day intervals while warm rainy weather conditions prevail.

FRUIT

CROP	DISEASES CONTROLLED	RATE/100 L	RATE/ha	WHP HARVEST (H) GRAZING (G)	CRITICAL COMMENTS
Bananas	Leaf spot (<i>Mycosphaerella musicola</i>) Leaf speckle Cordana leaf spot Fruit speckle Black pit	200 g plus spray oil (used according to its label directions)	1.25 to 1.60 kg plus 4 to 6 L Polyphase summer oil	Nil (H)	Apply when weather conditions favour disease outbreak. Good deleafing practices are essential for effective disease control. Remove infected leaf regularly. Use higher rates when spray coverage of Fortuna Globe 750 WG is compromised by aerial application conditions or terrain. Control of leaf spot may be improved with application by airblast sprayers from the ground. North Queensland, WA, NT: Repeat sprays at 2-4 week intervals. DO NOT extend spray intervals beyond 2 weeks until the plantation has been thoroughly de-leafed. DO NOT extend spray intervals beyond 2 weeks if conditions are favourable for disease, for example, wet and/or humid weather or if disease pressure is high. South Queensland, NSW: Repeat sprays at 2-3 week intervals during the warm, humid and/or wet months (i.e. December or January to May).

VEGETABLES

CROP	DISEASES CONTROLLED	RATE/100 L	RATE/ha	WHP HARVEST (H) GRAZING (G)	CRITICAL COMMENTS
Capsicum	Target spot	150 to 200 g	2 to 3 kg	14 days (H)	Apply when disease symptoms first appear and then repeat at 7 to 10 day intervals. Use the higher rate and shorter spray interval during weather conditions favourable to disease development.
Tomatoes	Early blight (target spot), Late blight (Irish blight), Anthracnose Leaf Mould (Fulvia fulva), Grey leaf spot, Phoma rot			7 days (H)	
Onion and Garlic	Downy mildew Purple blotch		2.2 to 3.5 kg		Apply when disease symptoms first appear and then repeat at 7 to 10 day intervals. Use the higher rate and shorter spray interval during weather conditions favourable to disease development.
Celery	Septoria leaf spot (leaf blight)		1.7 to 2.2 kg		
Carrots	Cercospora leaf spot Alternaria leaf spot				

VEGETABLES continued

CROP	DISEASES CONTROLLED	RATE/100 L	RATE/ha	WHP HARVEST (H) GRAZING (G)	CRITICAL COMMENTS
Lettuce	Anthraxnose Downy Mildew Septoria leaf spot	150 to 200 g	1.7 to 2.2 kg	14 days (H)	Apply when disease symptoms first appear and then repeat at 7 to 10 day intervals. Use the higher rate and shorter spray interval during weather conditions favourable to disease development.
Spinach Silver Beet Beetroot	Downy mildew Cercospora leaf spot				
Rhubarb	Downy Mildew Rust				
Potatoes	Early blight (Target spot), Late blight (Irish blight)			Not required	

COLE CROPS AND CUCURBITS

CROP	DISEASES CONTROLLED	RATE/100 L	RATE/ha	WHP HARVEST (H) GRAZING (G)	CRITICAL COMMENTS
Cabbage Cauliflower Brussels Sprouts Broccoli	Anthraxnose, Downy mildew, Alternaria spot, Ring spot	150 to 200 g	1.7 to 2.2 kg	7 days (H)	Apply when disease symptoms first appear and then repeat at 7 to 10 day intervals. Use the higher rate and shorter spray interval during weather conditions favourable to disease development.
Cucumber Cantaloupe Pumpkin Melons Zucchini Squash Pumpkin	Downy mildew Anthraxnose Gummy Stem blight Septoria spot				

PULSES GROWN FOR DRY BEANS

CROP	DISEASES CONTROLLED	RATE/ha	WHP HARVEST (H) GRAZING (G)	CRITICAL COMMENTS
Broad beans / Faba beans	Ascochyta blight Cercospora leaf spot Chocolate spot, Rust	1.0-2.2 kg	4 weeks (H) 7 days (G)	Spray when leaves, flowers or pods on most plants first show signs of infection and weather conditions are likely to remain humid or wet and favour disease development and repeat at 7-10 day intervals. If disease pressure is severe and weather conditions favour spread of the disease, repeat the spray treatment at 2-3 week intervals in order to protect new growth and developing pods. Use higher rate for dense crops and if disease is severe. Repeat more frequently if severe disease persists. For improved coverage add a registered surfactant at 10-20mL/ 100 L of spray.
Chickpeas	Ascochyta blight Botrytis Grey Mould			
Field peas	Ascochyta blight Black spot Botrytis Grey Mould Rust			
Lentils and Vetch	Ascochyta blight Botrytis Grey Mould Rust			
Lupins	Botrytis Grey Mould Anthraxnose			
Beans (<i>Phaseolus</i> spp.)	Rust, Anthracnose Ascochyta blight Angular leaf spot			

PEAS AND BEANS GROWN FOR FRESH CONSUMPTION

CROP	DISEASES CONTROLLED	RATE/ha	WHP HARVEST (H) GRAZING (G)	CRITICAL COMMENTS
Broad beans / Faba beans	Rust Cercospora leaf spot	1.7-2.2kg	7 days (H) 14 days (G)	Apply when disease symptoms first appear and then repeat at 7-10 day intervals. Use the higher rate and shorter spray interval during weather conditions favourable to disease development. Spray when leaves, flowers or pods show first sign of infection and weather conditions are likely to remain humid or wet and favour disease development.
	Chocolate spot Ascochyta blight (suppression only)	1.7-2.5 kg	7 days (H) 14 days (G)	If disease pressure is severe and weather conditions favour spread of the disease, repeat the spray treatment at 2-3 week intervals in order to protect new growth and developing pods. Use higher rate for dense crops and if disease is severe. Volume of application: for ground equipment apply 100 L/ha; for aircraft spray apply at least 30 L/ha.
Peas (<i>Pisum</i> spp)	Rust	1.7-2.2 kg		Apply when disease symptoms first appear and then repeat at 7-10 day intervals. Use the higher rate and shorter spray interval during weather conditions favourable to disease development. Spray when leaves, flowers or pods show first sign of infection and weather conditions are likely to remain humid or wet and favour disease development.
	Ascochyta leaf blight (suppression only)	1.7-2.5 kg		If disease pressure is severe and weather conditions favour spread of the disease, repeat the spray treatment at 2-3 week intervals in order to protect new growth and developing pods. Use higher rate for dense crops and if disease is severe. Volume of application: for ground equipment apply 100 L/ha; for aircraft spray apply at least 30 L/ha.
Green Beans (<i>Phaseolus</i> spp.)	Rust Anthracnose Angular leaf spot	1.7 to 2.2 kg		Apply when disease symptoms first appear and then repeat at 7-10 day intervals. Use the higher rate and shorter spray interval during weather conditions favourable to disease development. Spray when leaves, flowers or pods show first sign of infection and weather conditions are likely to remain humid or wet and favour disease development.

TURF

CROP	DISEASES CONTROLLED	RATE/100 L	CRITICAL COMMENTS
Turf	Fusarium Rhizoctonia Helminthosporium	200 to 250 g / 100 sq. metres	Apply as a protective spray schedule at 7-day intervals throughout the susceptible season. Use sufficient water for adequate coverage. Do not feed grass clippings from treated areas to poultry or animals.

ORNAMENTALS

CROP	DISEASES CONTROLLED	RATE/100 L	CRITICAL COMMENTS
Roses	Black spot	150 to 200 g	Apply when disease symptoms first appear and then repeat at 7-10 day intervals. Use the higher rate and shorter spray interval during weather conditions favourable to disease development. Caution: This product may be Phytotoxic to some ornamental species (in particular some varieties of ferns). It is recommended to test a few plants before widespread use.
Gladioli Chrysanth-emums	Grey mould (<i>Botrytis</i>) Septoria leaf spot Rust		
Carnations	Rust Alternaria leaf spot		
Flowers Shrubs	Septoria leaf spot		
Ferns	Botrytis leaf spot	75 to 100 g	

TREE AND VINE CROPS

RATE			CRITICAL COMMENTS
In the following table, all rates are given for dilute spraying. For concentrate spraying, refer to the Application Section.			For all uses in the table: Apply by dilute or concentration spraying equipment. Apply the same amount of product to the target crop whether applying this product by dilute or concentrate spraying methods.
CROP	DISEASES CONTROLLED	RATE/100 L	
POME FRUITS Apples Pears	Pear scab Apple scab (Black spot) Target spot Bitter rot Sooty blotch Fly speck Rite fruit spot	150 to 200 g	14 days (H)
			Apply at 7 to 14 day intervals following a copper spray at green tip. Use the higher rate and shorter spray interval during prolonged wet conditions. This product may be harmful to <i>Typhlodromus pyri</i> and its use in orchards where integrated control is practiced should be avoided.

TREE AND VINE CROPS continued

RATE				CRITICAL COMMENTS
In the following table, all rates are given for dilute spraying. For concentrate spraying, refer to the Application Section.				For all uses in the table: Apply by dilute or concentration spraying equipment. Apply the same amount of product to the target crop whether applying this product by dilute or concentrate spraying methods.
CROP	DISEASES CONTROLLED	RATE/100 L	WHP HARVEST (H) GRAZING (G)	
STONE FRUIT Almonds Peaches Apricots Plums (except early plums such as Wilson & Wickson) Nectarines Cherries	Rust Shot Hole Freckle Brown Rot	150 to 200 g	14 days (H)	Apply at early bloom (1-10%), then repeat at mid to full bloom (50-100%); at petal fall, and at shuck fall. Continue with a protective spray programme at two-week intervals. Caution: This product may be Phytotoxic to Wilson and Wickson Plums. It is advisable to test any new plum variety on a small scale with Fortuna Globe 750 WG before applying to the whole crop.
Passion fruit	Septoria spot Brown spot (<i>Alternaria passiflorae</i>)	150 g	1 day (H)	Apply at 10 to 14 day intervals from October to May and every 21 to 28 days for remainder of year.
	Anthracnose	200 g plus 600 mL miscible summer oil		Spray at weekly intervals during flowering and then fortnightly until near harvest.
Pawpaws (Papaya)	Black spot	200 g	1 day (H)	Apply Fortuna Globe 750 WG plus a wetting agent at 10-14 day intervals during periods conducive to disease.
Custard apples	Pseudocercospora fruit spot			Do not apply during flowering. Apply as 3-4 week intervals from first fruit set until harvest. Apply at 2 week intervals in wet, cloudy weather.
Mangoes	Anthracnose (<i>Collectotrichum gloeosporioides</i>)			Apply at weekly intervals during flowering and then monthly intervals until harvest.
Grapevines	Blackspot Downy mildew	150 to 200 g	30 days (H)	For Black spot control apply at budburst and then repeat 10 to 14 days later. If Downy mildew is expected, continue spray programme at 10 to 14 day intervals until the threat of disease has passed. Use the shorter spray interval during weather conditions favourable to disease development.
	Phomopsis Cane and Leaf spot			Apply at budburst and then repeat 7 to 10 days later.
Citrus	Black spot Citrus rust mite Brown citrus mite	200 g plus spray oil (used according to directions)	Not required	Add an adjuvant (spreader/sticker) according to the adjuvant label recommendation per 100 L. Choose a spray oil recommended for use on citrus. On heavy soil: Spray at eight and at fifteen weeks following a Bordeaux (5-5-100) spray at petal fall. On light soil: Spray at six and at thirteen weeks following a Bordeaux (3.5- 3.5-100) spray at petal fall. Queensland: Spray at six and twelve weeks after a copper spray at ½ to ¾ petal fall. Apply at first sign of mite activity on fruit, usually between December and May.
	Citrus bud mite			Apply when blossom malformation is observed, usually February. Choose a spray oil recommended for use on citrus.

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

WITHHOLDING PERIOD PRIOR TO HARVEST:

BANANAS – NOT REQUIRED WHEN USED AS DIRECTED.

MANGOES, PASSION FRUIT, CUSTARD APPLES – DO NOT HARVEST FOR 1 DAY AFTER APPLICATION.

PAWPAWS (PAPAYA) – DO NOT HARVEST FOR 1 DAY AFTER APPLICATION. PAPAYA LEAVES MUST NOT BE MADE AVAILABLE FOR HUMAN CONSUMPTION.

BULB VEGETABLES, CELERY, CARROTS, COLE CROPS, CUCURBITS, TOMATOES, SOYBEANS, GREEN BEANS – DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.

CAPSICUMS, LETTUCE, SPINACH, SILVER BEET, BEETROOT, RHUBARB, POME AND STONE FRUIT, NECTARINES, ALMONDS, PEANUTS, FENNEL – DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION.

GRAPEVINES – DO NOT HARVEST FOR 30 DAYS AFTER APPLICATION.

BROAD BEANS, FABA BEANS, PEAS:

- HARVEST for FRESH CONSUMPTION – DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.

- HARVEST for DRY CONSUMPTION – DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION.

- CHICKPEAS, FIELD PEAS, LUPINS LENTILS, VETCH – DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION.

- POPPIES – DO NOT HARVEST FOR 7 WEEKS AFTER APPLICATION.

CITRUS, POTATOES, TOBACCO – NOT REQUIRED WHEN USED AS DIRECTED.

WITHHOLDING PERIOD PRIOR TO GRAZING:

SOYBEANS, PEANUTS – DO NOT GRAZE OR CUT FOR STOCKFEED FOR 14 DAYS AFTER APPLICATION.

BEANS, FIELD PEAS, CHICKPEAS, LENTILS, LUPINS, VETCH – DO NOT GRAZE OR CUT FOR STOCKFEED FOR 14 DAYS AFTER APPLICATION.

PAWPAWS (PAPAYA) – DO NOT USE OR SUPPLY TREATED PAPAYA INCLUDING LEAVES FOR STOCK FOOD.

COTTON – DO NOT GRAZE LIVESTOCK ON TREATED CROPS OR FEED TREATED CROPS, STUBBLE AND/OR TRASH (INCLUDING COTTON GIN TRASH) TO LIVESTOCK.

SUNFLOWERS – DO NOT GRAZE LIVESTOCK ON TREATED CROPS OR FEED TREATED CROPS AND/OR TRASH TO LIVESTOCK.

APVMA approval number: 67494/109664

Additional statements as required by Safe Work Australia in accordance with the Globally Harmonized System of Classification and Labelling (GHS)

Warning

May cause an allergic skin reaction. Suspected of damaging fertility or the unborn child. Very toxic to aquatic life. Contaminated work clothing should not be allowed out of the workplace. Avoid release to the environment. Wear protective gloves/protective clothing/eye protection/face protection. IF ON SKIN: Wash with plenty of soap and water. Wash contaminated clothing before reuse. Collect spillage.

In a Transport Emergency
Dial 000
Police or Fire Brigade

Specialist Advice In Emergency Only
1800 033 111
All Hours - Australia Wide